

KKM Property and Fixed Point Theorems*

Tong-Huei Chang

*Department of Business Administration, Shih Chien College,
Taipei, Taiwan, Republic of China*

Metadata, citation and similar papers at core.ac.uk

Chn-Lin Yen

*Department of Mathematics, National Taiwan Normal University,
Taipei, Taiwan, Republic of China*

Submitted by Jean Mawhin

Received May 15, 1995

1. INTRODUCTION

Recently, the famous Knaster–Kuratowski–Mazurkiewicz theorem [9] and Fan's lemma [5] have widely been used as very versatile tools in modern nonlinear analysis (see, for example, [1–3, 8, 10, 18, 19]).

It is well known that if T is a KKM mapping from a convex subset X of a topological vector space E into 2^E with closed values then the family $\{Tx: x \in X\}$ has the finite intersection property. Many authors had generalized the above result to the following form.

Let X be a convex space, Y a topological space, and let S, T be two set-valued mappings from X into 2^Y such that for any finite subset $\{x_1, x_2, \dots, x_n\}$ of X , $T(\text{co}\{x_1, \dots, x_n\}) \subset \bigcup_{i=1}^n Sx_i$, then the family $\{\bar{S}x: x \in X\}$ has the finite intersection property under some assumptions of T (for example, one may see [11–14]).

In this paper, we shall use the above results to define a family of set-valued mappings from a set X into a topological space Y . We get some fixed point theorems and coincidence theorems which extend the results of

* This work was partially supported by the National Science Council of the Republic of China under Contract NSC 84-2121-M-003-008.

[7, 14, 15]. We also get a generalized Fan's matching theorem and a generalized KKM theorem. As applications, we use these results to study the existence problem of solutions for some generalized variational inequalities.

2. PRELIMINARIES

Let X and Y be two sets, and let $T: X \rightarrow 2^Y$ be a set-valued mapping where $2^Y = \{B \subset Y: B \text{ is nonempty}\}$, we shall use the following notions in the sequel:

- (i) $TA = \bigcup_{x \in A} Tx$ for $A \subset X$;
- (ii) $T^{-1}y = \{x \in X: y \in Tx\}$ for $y \in Y$;
- (iii) $T^{-1}B = \bigcup_{y \in B} T^{-1}y$ for $B \subset Y$.

For the case that X and Y are two topological spaces,

(iv) T is said to be upper semi-continuous (u.s.c.) if for each closed subset B of Y , $T^{-1}B$ is closed in X ;

(v) T is said to be closed if the graph $\Gamma_T = \{(x, y) \in X \times Y: y \in Tx\}$ is a closed subset of $X \times Y$; and

(vi) T is said to be compact if the image TX of X under T is contained in a compact subset of Y .

Assume that X is a convex subset of a linear space and Y is a topological space. If $S, T: X \rightarrow 2^Y$ are two set-valued mappings such that $T(\text{co } A) \subset SA$ for each finite subset A of X , then we call S a generalized KKM mapping w.r.t. T , where $\text{co } A$ denotes the convex hull of A . Let $T: X \rightarrow 2^Y$ be a set-valued mapping such that if $S: X \rightarrow 2^Y$ is a generalized KKM mapping w.r.t. T then the family $\{\overline{Sx}: x \in X\}$ has the finite intersection property (where \overline{Sx} denotes the closure of Sx), then we say that T has the KKM property. Denote $\text{KKM}(X, Y) = \{T: X \rightarrow 2^Y | T \text{ has the KKM property}\}$.

Remark. (1) Generalized KKM mappings were first introduced by Park [13], and followed by some others.

(2) A closed-valued generalized KKM set-valued mapping w.r.t. some set-valued mapping may not have the finite intersection property (f.i.p.).

For example, if $X = [0, 1]$ with the usual topology and $S, T: X \rightarrow 2^X$ are defined as

$$Sx = Tx = \begin{cases} \{0\}, & x \in [0, \frac{1}{2}); \\ \{0, 1\}, & x = \frac{1}{2}; \\ \{1\}, & x \in (\frac{1}{2}, 1], \end{cases}$$

then

- (i) Sx is closed (compact) for each $x \in X$;
- (ii) T is compact, closed (hence u.s.c.); and
- (iii) $T(\text{co } A) \subset SA$ for each finite subset A of X ,

however, the family $\{Sx: x \in X\}$ has no f.i.p.

A convex space S is a convex set (in a linear space) with any topology that induces the Euclidean topology on the convex hull of its finite subset. For each finite subset A of X , the convex hull of A is called a polytope of X . Let Y be a topological space, a subset B of Y is compactly closed if for any compact subset K of Y , $B \cap K$ is closed in K .

Let \mathbf{F} be a class of set-valued mappings. Denote

$$\mathbf{F}(X, Y) = \{T: X \rightarrow 2^Y | T \in \mathbf{F}\}$$

$\mathbf{F}_c = \{T_n T_{n-1} \dots T_1 | T_i \in \mathbf{F}, i = 1, 2, \dots, n \text{ for some } n\}$, that is, the set of finite composites of mappings in \mathbf{F} .

If X and Y are two topological spaces, we note

$$\mathbf{C}(X, Y) = \{s: X \rightarrow Y | s \text{ is a single-valued continuous function}\};$$

$\mathbf{K}(X, Y) = \{T: X \rightarrow 2^Y | T \text{ is u.s.c. with compact convex values and } Y \text{ is convex}\};$

$\mathbf{V}(X, Y) = \{T: X \rightarrow 2^Y | T \text{ is u.s.c. with compact acyclic values}\}$ (see [15]); and

$\mathbf{A}(X, Y) = \{T: X \rightarrow 2^Y | T \text{ is u.s.c. and approachable with compact values}\}$ (see [3]).

The following abstract class \mathfrak{A} of set-valued mappings is introduced by Park [14].

A class \mathfrak{A} of set-valued mappings is one satisfying the following:

- (i) \mathfrak{A} contains the class \mathbf{C} of single-valued continuous functions;
- (ii) each $T \in \mathfrak{A}_c$ is u.s.c. with compact values; and
- (iii) for any polytope Δ , each $T \in \mathfrak{A}_c(\Delta, \Delta)$ has a fixed point.

Denote $\mathfrak{A}_c^*(X, Y) = \{T: X \rightarrow 2^Y | \text{for any compact subset } K \text{ of } X, \text{ there is a } F \in \mathfrak{A}_c(K, Y) \text{ such that } Fx \subset Tx \text{ for each } x \in K\}$.

3. MAIN RESULTS

In the sequel, all topological spaces are supposed to satisfy the Hausdorff separation property.

We first prove the following lemma:

LEMMA 1. *Let X be a convex subset of a topological vector space E , V a symmetric convex open neighborhood of 0 in E . If $T \in \text{KKM}(X, X)$ is compact, then there is $x_v \in X$ such that $(x_v + V) \cap Tx_v \neq \emptyset$.*

Proof. Suppose that $(x + V) \cap Tx = \emptyset$ for each $x \in X$. Let $X_1 = \overline{TX}$ then X_1 is a compact subset of X since T is compact. Define the mapping $S: X \rightarrow 2^X$ by

$$Sx = X_1 \setminus (x + \frac{1}{4}V) \quad \text{for each } x \in X.$$

We have

- (a) Sx is closed in X for each $x \in X$; and
- (b) S is a generalized KKM mapping w.r.t. T .

To prove (b), it suffices to show that for any finite subset $A = \{x_1, x_2, \dots, x_n\}$ of X , $T(\text{co } A) \subset SA$.

It is trivial for the case that $n = 1$. We now consider $n \geq 2$.

Case (i). If there is some i , $2 \leq i \leq n$, such that $(x_i + \frac{1}{4}V) \cap (x_1 + \frac{1}{4}V) = \emptyset$, then $T(\text{co } A) \subset X_1 = Sx_i \cup Sx_1 \subset SA$.

Case (ii). Suppose that $(x_i + \frac{1}{4}V) \cap (x_1 + \frac{1}{4}V) \neq \emptyset$ for each $2 \leq i \leq n$. We have $x_1 \in x_i + \frac{1}{2}V$, for each $2 \leq i \leq n$. Hence $x_1 \in x + \frac{1}{2}V$ for each $x \in \text{co } A$. This implies $x_1 + \frac{1}{2}V \subset \bigcap_{x \in \text{co } A} (x + V)$ and so $T(\text{co } A) \subset Sx_1 \subset SA$.

Since $T \in \text{KKM}(X, X)$, hence the family $\{Sx: x \in X\}$ has f.i.p. and hence $\bigcap_{x \in X} Sx \neq \emptyset$. Let $x_0 \in \bigcap_{x \in X} Sx$, then $x_0 \in Sx_0$. This contradicts the definition of S .

Using Lemma 1, we get our main theorem:

THEOREM 2. *Let X be a convex subset of a locally convex space E , and $T \in \text{KKM}(X, X)$. If T is compact and closed, then T has a fixed point in X .*

Proof. Suppose $\{V_\alpha: \alpha \in \Lambda\}$ is a local basis of E such that V_α is symmetric, open, and convex for each $\alpha \in \Lambda$.

By Lemma 1, for each $\alpha \in \Lambda$, there exists $x_\alpha \in X$ such that $Tx_\alpha \cap (x_\alpha + V_\alpha) \neq \emptyset$. That is, there exists $y_\alpha \in Tx_\alpha$ such that $y_\alpha \in x_\alpha + V_\alpha$ for each $\alpha \in \Lambda$. Since T is compact, without loss of generality, we may assume that y_α converges to some \hat{y} in Y , then x_α also converges to \hat{y} . Since T is closed, we have $\hat{y} \in T\hat{y}$.

We now list some properties of the KKM family.

PROPOSITION 3. *Let X be a convex subset of a linear space, and let Y, Z be two topological spaces. Then*

- (i) *$T \in \text{KKM}(X, Y)$ if and only if $T|_{\Delta} \in \text{KKM}(\Delta, Y)$ for each polytope Δ in X ;*
- (ii) *if $T \in \text{KKM}(X, Y)$ and $f \in \mathbf{C}(Y, Z)$, then $fT \in \text{KKM}(X, Z)$;*
- (iii) *if Y is a normal space, X a convex space, Δ a polytope in X , and if $T: \Delta \rightarrow 2^Y$ is a set-valued mapping such that for each $f \in \mathbf{C}(Y, \Delta)$, fT has a fixed point in Δ , then $T \in \text{KKM}(\Delta, Y)$; and*
- (iv) *if X is a convex space, then $\mathfrak{K}_c^\kappa(X, Y) \subset \text{KKM}(X, Y)$.*

Proof. Part (i) is obvious. To prove (ii), let $S: X \rightarrow 2^Z$ be a generalized KKM mapping w.r.t. fT such that Sx is closed for each $x \in X$. Then, for any finite subset $\{x_1, x_2, \dots, x_n\}$ of X , since S is a generalized KKM mapping w.r.t. fT , we have

$$fT(\text{co}\{x_1, x_2, \dots, x_n\}) \subset \bigcup_{i=1}^n Sx_i,$$

and hence $T(\text{co}\{x_1, x_2, \dots, x_n\}) \subset \bigcup_{i=1}^n f^{-1}Sx_i$. Therefore, $f^{-1}S$ is a generalized KKM mapping w.r.t. T . Since $T \in \text{KKM}(X, Y)$, hence the family $\{f^{-1}Sx: x \in X\}$ has f.i.p., and hence the family $\{Sx: x \in X\}$ has f.i.p. This shows that $fT \in \text{KKM}(X, Z)$ and we complete the proof of (ii).

The proof of (iii) is similar to the proof of [11, Lemma 2] and we omit it. Part (iv) is a direct consequence of [14, Corollary 2].

Remark. (1) Since \mathbf{C} , \mathbf{K} , and \mathbf{V} are examples of \mathfrak{K} , hence, by Proposition 3(iv), $\mathbf{K}_c \subset \text{KKM}$ and $\mathbf{V}_c \subset \text{KKM}$.

(2) The following example shows that $\text{KKM} \not\supseteq \mathfrak{K}_c^\kappa$, hence Theorem 2 properly extends Theorems 3(iii) of [14].

EXAMPLE. Let $X = Y = [0, 1]$ with the usual topology, and let $T: X \rightarrow 2^Y$ be defined as

$$Tx = \begin{cases} \left\{ \left| \sin \frac{1}{x} \right| \right\}, & x \in (0, 1]; \\ \{0\}, & x = 0. \end{cases}$$

Then $T \in \text{KKM}(X, Y)$, however $T \notin \mathfrak{K}_c^\kappa(X, Y)$.

By Proposition 3 and Theorem 2, we get the following coincidence theorem which extends a result of Park, Singh and Watson [15, Theorem 1] and the main result of Granas and Liu [7].

THEOREM 4. *Let X be a convex space, and let $T, G: X \rightarrow 2^Y$ be two set-valued mappings satisfying:*

- (i) $T \in \text{KKM}(X, Y)$ is compact and closed;
- (ii) for each $y \in GX$, $G^{-1}y$ is convex; and
- (iii) $\{\text{int } Gx: x \in X\}$ covers \overline{TX} .

Then there exists $x_0 \in X$ such that $Tx_0 \cap Gx_0 \neq \emptyset$.

Proof. Since T is compact and $\overline{TX} \subset \bigcup_{x \in X} \text{int } Gx$, hence there exists a finite subset $\{x_1, x_2, \dots, x_n\}$ of X such that $\overline{TX} \subset \bigcup_{i=1}^n \text{int } Gx_i$. Let $\Delta = \text{co}\{x_1, x_2, \dots, x_n\}$, and let $\{\alpha_i\}_{i=1}^n$ be the partition of the unity subordinated to $\{\text{int } Gx_i: i = 1, 2, \dots, n\}$. Define $f: \overline{TX} \rightarrow \Delta$ by

$$f(y) = \sum_{i=1}^n \alpha_i(y)x_i \quad \text{for each } y \in \overline{TX}$$

then $f \in \mathbf{C}(\overline{TX}, \Delta)$. By Proposition 3(i), $T|_{\Delta} \in \text{KKM}(\Delta, \overline{TX})$, and by Proposition 3(ii) and Theorem 2, fT has a fixed point in Δ , that is, there exists $x_0 \in \Delta$ such that $x_0 \in fTx_0$. Let $y_0 \in Tx_0$ such that $x_0 = f(y_0)$, and let $I(y_0) = \{i \in \{1, 2, \dots, n\}: \alpha_i(y_0) > 0\}$. Then $x_0 = \sum_{i \in I(y_0)} \alpha_i(y_0)x_i$. If $\alpha_i(y_0) > 0$, then $y_0 \in \text{int } Gx_i \subset Gx_i$, and then $x_i \in G^{-1}y_0$. So $x_0 \in \text{co}\{x_i: i \in I(y_0)\} \subset \text{co}(G^{-1}y_0) = G^{-1}y_0$, since $G^{-1}y_0$ is convex. This implies $x_0 \in G^{-1}y_0$ and so $y_0 \in Gx_0$. Thus $y_0 \in Tx_0 \cap Gx_0$ and we complete our proof.

We now use a result of Ben-El-Mechaiekh and Deguire to get some fixed point theorems and a generalized Fan's matching theorem.

PROPOSITION 5. *Let X be a convex subset of a locally convex space E , Y a compact subset of a topological vector space F , and let $T \in \text{KKM}(X, Y)$ be closed. Then for any $G \in \mathbf{A}_c(Y, X)$, TG has a fixed point in Y (and hence GT has a fixed point in X).*

Proof. Since $T \in \text{KKM}(X, Y)$, hence for any $f \in \mathbf{C}(Y, X)$, by Proposition 3(ii), $fT \in \text{KKM}(X, X)$, and hence, by Theorem 2, fT has fixed point in X . That is, $\Gamma_f \cap \Gamma_{T^{-1}} \neq \emptyset$. So, by [3, Corollary 7.5], we have $\Gamma_G \cap \Gamma_{T^{-1}} \neq \emptyset$. That is, TG has a fixed point in Y .

COROLLARY 6. *Let X and Y be defined as in Proposition 5, and let Z be a subset of a topological vector space L . If $T \in \text{KKM}(X, Y)$ is closed, then for each $G \in \mathbf{A}_c(Y, Z)$, $GT \in \text{KKM}(X, Z)$.*

Proof. By Proposition 3(i), it suffices to show that $GT|_{\Delta} \in \text{KKM}(\Delta, Z)$, for each polytope Δ in X . Since for any $f \in \mathbf{C}(Z, \Delta)$, $fG \in \mathbf{A}_c(Y, \Delta)$, so,

by Proposition 5, $fGT|_{\Delta}$ has a fixed point in Δ . Thus, by Proposition 3(iii), $GT|_{\Delta} \in \text{KKM}(\Delta, Z)$.

Remark. It also can be shown, under some suitable modification, that for any $T \in \text{KKM}(X, Z)$ and $G \in \mathbf{A}_c(Y, X)$, we have $TG \in \text{KKM}(Y, Z)$.

Using the above result, we have the following theorem, which is a generalized Fan's matching theorem:

THEOREM 7. *Let X be a compact convex subset of a locally convex space, and let $\{A_{\nu}: \nu \in I\}$ be a finite family of closed subsets of X indexed by a set I , such that $\bigcup_{\nu \in I} A_{\nu} = X$. If $T \in \text{KKM}(X, X)$ is compact and closed, then for any family $\{x_{\nu}\}_{\nu \in I}$ of points of X , indexed by the same set I , there exists a nonempty subset J of I such that*

$$T(\text{co}\{x_{\nu}: \nu \in J\}) \cap \left(\bigcap_{\nu \in J} A_{\nu} \right) \neq \emptyset.$$

Proof. For each $x \in X$, let $I(x) = \{\nu \in I: x \in A_{\nu}\}$, then $I(x) \neq \emptyset$ for each $x \in X$ since $\{A_{\nu}\}_{\nu \in I}$ covers X .

Define $F: X \rightarrow 2^X$ by

$$Fx = \text{co}\{x_{\nu}: \nu \in I(x)\} \quad \text{for each } x \in X.$$

It is clear that Fx is a nonempty compact convex subset of X . For each $x \in X$, let $U(x) = X \setminus \bigcup\{A_{\nu}: \nu \notin I(x)\}$, then $U(x)$ is an open neighborhood of x . If $z \in U(x)$, then $Fz \subset Fx$. This shows that F is u.s.c., hence $F \in \mathbf{K}(X, X)$, and hence, by Proposition 4.1 of [3], $F \in \mathbf{A}(X, X)$. Now, by Proposition 5, there exists $x_0 \in X$ such that $x_0 \in TFx_0 = T(\text{co}\{x_{\nu}: \nu \in I(x_0)\})$. So,

$$x_0 \in T(\text{co}\{x_{\nu}: \nu \in I(x_0)\}) \cap \left(\bigcap_{\nu \in I(x_0)} A_{\nu} \right)$$

which completes the proof.

Remark. Fan [6, Theorem 1] proved Theorem 7 for the case that $T = i_x$ (the identity mapping of X).

4. APPLICATIONS

In this section, we shall use the results in Section 3 and the following generalized KKM theorem to get some generalized variational inequalities.

THEOREM 8. *Let X be a convex space, Y a topological space, let $S: X \rightarrow 2^Y$ be a set-valued mapping, and let $T \in \text{KKM}(X, Y)$ such that*

- (i) *for each compact subset X' of X , $\overline{TX'}$ is compact in Y ;*
- (ii) *for each $x \in X$, Sx is compactly closed in Y ;*
- (iii) *for any finite subset A of X , $T(\text{co } A) \subset SA$; and*
- (iv) *there exist a nonempty compact convex subset X_0 of X and a compact subset K of Y such that $\bigcap_{x \in X_0} Sx \subset K$.*

Then $\bigcap_{x \in X} Sx \neq \emptyset$.

Proof. Suppose that $\bigcap_{x \in X} Sx = \emptyset$, then $\bigcup_{x \in X} S^c x = Y$, and $\bigcup_{x \in X} (S^c x \cap K) = K$. Since K is compact and $S^c x$ is compactly open, there exists a finite subset $\{x_1, x_2, \dots, x_n\}$ of X such that $\bigcup_{i=1}^n (S^c x_i \cap K) = K$. By assumption (iv), $\bigcap_{x \in X_0} Sx \subset K$, we have $\bigcup_{x \in X_0} S^c x \supset K^c$. Now, let $X_1 = \text{co}(X_0 \cup \{x_1, x_2, \dots, x_n\})$ then X_1 is a compact convex set and $\bigcup_{x \in X_1} S^c x = Y$. Hence $\bigcap_{x \in X_1} Sx = \emptyset$.

By (i), $\overline{TX_1}$ is compact in Y . Define $F: X_1 \rightarrow 2^Y$ by

$$Fx = Sx \cap \overline{TX_1} \quad \text{for each } x \in X_1.$$

Then

- (a) Fx is closed (compact) in Y for each $x \in X_1$, and
- (b) $T(\text{co } A) \subset FA$, for any finite subset A of X_1 .

Since $T|_{X_1} \in \text{KKM}(X, Y)$, the family $\{Fx: x \in X_1\}$ has f.i.p., hence $\bigcap_{x \in X_1} Fx \neq \emptyset$, and hence $\bigcap_{x \in X_1} Sx \neq \emptyset$. This is a contradiction, and we complete our proof.

Remark. Condition (i) of Theorem 8 is weaker than the assumption that T is compact.

As a consequence of the above theorem, we have the following theorem, which is generalized form of [16, Theorem 3].

THEOREM 9. *Let X and Y be defined as in Theorem 8, and let $T \in \text{KKM}(X, Y)$ such that for each compact subset X' of X , $\overline{TX'}$ is compact in Y . If $\psi, \varphi: X \times Y \rightarrow \mathbb{R}$ are two real-valued functions satisfying:*

- (i) $\psi(x, y) \leq 0$, for each $(x, y) \in \Gamma_T$;
- (ii) for fixed $x \in X$, the mapping $y \rightarrow \varphi(x, y)$ is lower semicontinuous (l.s.c.) on A for each compact subset A of Y ;
- (iii) for fixed $y \in Y$, the set $\{x \in X: \psi(x, y) > 0\}$ contains the convex hull of the set $\{x \in X: \varphi(x, y) > 0\}$; and

(iv) *there exist a nonempty compact convex subset X_0 of X , and a compact subset K of Y such that for each $y \in Y \setminus K$, there is $x \in X_0$ such that $\varphi(x, y) > 0$,*

then there exists $\hat{y} \in K$ such that $\varphi(x, \hat{y}) \leq 0$ for each $x \in X$.

Proof. Define $F, S: X \rightarrow 2^Y$ by

$$Fx = \{y \in Y: \psi(x, y) \leq 0\}$$

and

$$Sx = \{y \in Y: \varphi(x, y) \leq 0\} \quad \text{for each } x \in X.$$

By assumption (i), we have $\Gamma_T \subset \Gamma_F$, and by assumption (ii), Sx is compactly closed for each $x \in X$. Condition (iii) implies that for each finite subset A of X , $F(\text{co } A) \subset SA$, and then $T(\text{co } A) \subset SA$.

Condition (iv) is equivalent to Theorem 8(iv). So, all the conditions in Theorem 8 are satisfied, and so $\bigcap_{x \in X} Sx \neq \emptyset$. Take $\hat{y} \in \bigcap_{x \in X} Sx$, and we have $\varphi(x, \hat{y}) \leq 0$ for each $x \in X$.

For the case where T is a single-valued mapping, we have the following corollary:

COROLLARY 10. *Let X and Y be defined as in Theorem 8, and let $s \in \mathbf{C}(X, Y)$. If $\psi, \varphi: X \times Y \rightarrow R$ are two real-valued functions such that*

(i) *$\psi(x, sx) \leq 0$, for each $x \in X$;*

(ii) *for fixed $x \in X$, the mapping $y \rightarrow \varphi(x, y)$ is l.s.c. on A for each compact subset A of Y ;*

(iii) *for fixed $y \in Y$, the set $\{x \in X: \psi(x, y) > 0\}$ contains the convex hull of the set $\{x \in X: \varphi(x, y) > 0\}$; and*

(iv) *there exist a nonempty compact convex subset X_0 of X , and a compact subset K of Y such that for each $Y \in Y \setminus K$ there is $x \in X$ such that $\varphi(x, y) > 0$,*

then there exist $\hat{y} \in K$ such that $\varphi(x, \hat{y}) \leq 0$ for each $x \in X$.

Remark. For the case $X = Y$ and $s = i_X$ (the identity mapping of X), Corollary 10 reduces to [16, Theorem 3].

Let X be a convex subset of a linear space, Y a set, and let $\psi, \varphi: X \times Y \rightarrow R$ be two real-valued functions. For any $y \in Y$, ψ is said to be φ -quasiconcave in x if for any finite subset $A = \{x_1, x_2, \dots, x_n\}$ of X , we have

$$\psi(x, y) \geq \min_{1 \leq i \leq n} \varphi(x_i, y) \quad \text{for all } x \in \text{co } A.$$

It is clear that if $\varphi(x, y) \leq \psi(x, y)$ for each $(x, y) \in X \times Y$, and if for each $y \in Y$, the mapping $x \rightarrow \varphi(x, y)$ is quasiconcave (or the mapping $x \rightarrow \psi(x, y)$ is quasiconcave) then ψ is φ -quasiconcave in x .

By the above definition, we have the following theorem concerning other inequalities.

THEOREM 11. *Let X be a convex space, Y a topological space, and let $T \in \text{KKM}(X, Y)$ be compact. If $\psi, \varphi: X \times Y \rightarrow R$ are two real-valued functions satisfying that*

- (i) *for each $x \in X$, the mapping $y \rightarrow \varphi(x, y)$ is l.s.c. on Y ; and*
- (ii) *ψ is φ -quasiconcave in x for each $y \in Y$,*

then for each $\lambda \in R$ one of the following properties holds:

- (a) *there exists $\hat{y} \in Y$ such that*

$$\varphi(x, \hat{y}) \leq \lambda \quad \text{for all } x \in X;$$

- (b) *for some $(x_0, y_0) \in \Gamma_T$ we have*

$$\psi(x_0, y_0) > \lambda.$$

Proof. Let $\lambda \in R$. Since T is compact, hence \overline{TX} is compact in Y . Define $F, S: X \rightarrow 2^Y$ by

$$Fx = \{y \in \overline{TX}: \psi(x, y) \leq \lambda\}$$

and

$$Sx = \{y \in \overline{TX}: \varphi(x, y) \leq \lambda\} \quad \text{for each } x \in X.$$

Suppose the conclusion (b) fails, then for each $(x, y) \in \Gamma_T$, $\psi(x, y) \leq \lambda$ and then $\Gamma_T \subset \Gamma_F$. Now, by (ii), we have $F(\text{co } A) \subset SA$ for each finite subset A of X . So, for any finite subset A of X , $T(\text{co } A) \subset SA$. Since $T \in \text{KKM}(X, Y)$ and, by (i), Sx is closed (compact), the family $\{Sx: x \in X\}$ has f.i.p. Hence $\bigcap_{x \in X} Sx \neq \emptyset$. Take $\hat{y} = \bigcap_{x \in X} Sx$, and we have

$$\varphi(x, \hat{y}) \leq \lambda \quad \text{for each } x \in X.$$

For the case that T is a single-valued function, we have the following corollary:

COROLLARY 12. *Let X be a convex space, Y a topological space, and let $s \in \mathbf{C}(X, Y)$ be compact. If $\psi, \varphi: X \times Y \rightarrow R$ are two real-valued functions*

satisfying conditions (i) and (ii) of Theorem 11, then for each $\lambda \in R$, one of the following properties holds:

(a) there exists $\hat{y} \in Y$ such that

$$\varphi(x, \hat{y}) \leq \lambda \quad \text{for all } x \in X;$$

(b) for some $x_0 \in X$ we have $\psi(x_0, sx_0) > \lambda$.

Remark. (1) Theorems 5.1 and 5.2 of [7] are special cases of Theorem 11 and Corollary 12, respectively.

(2) The conclusion of Theorem 11 implies the inequality

$$\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \sup_{(x, y) \in \Gamma_T} \psi(x, y),$$

and the conclusion of Corollary 12 implies the inequality

$$\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \sup_{x \in X} \psi(x, sx),$$

which are the generalized forms of [18, Theorem 1].

We now present another result by using the KKM property directly.

THEOREM 13. *Let X be a convex space, Y a compact space. Let $\psi, \varphi: X \times Y \rightarrow R$ be two real-valued functions satisfying conditions (i) and (ii) of Theorem 11, and let $\lambda \in R$. If for any polytope Δ in X and for $s \in \mathbf{C}(Y, \Delta)$, there exists $y \in Y$ (depends on s) such that $\psi(sy, y) \leq \lambda$, then there exists $\hat{y} \in Y$ such that*

$$\varphi(x, \hat{y}) \leq \lambda \quad \text{for all } x \in X.$$

Proof. Define $T, S: X \rightarrow 2^Y$ by

$$Tx = \{y \in Y: \psi(x, y) \leq \lambda\}$$

and

$$Sx = \{y \in Y: \varphi(x, y) \leq \lambda\} \quad \text{for each } x \in X.$$

By assumption, for any polytope Δ in X and for $s \in \mathbf{C}(Y, \Delta)$, $sT|_{\Delta}$ has a fixed point in Δ . Hence, by Proposition 3(iii), $T \in \text{KKM}(X, Y)$. The rest of the proof is similar to the proof of Theorem 11 and we omit it.

ACKNOWLEDGMENT

The authors are grateful to the referee for his valuable suggestions.

REFERENCES

1. J. P. Aubin, "Mathematical Methods of Game and Economic Theory," North-Holland, New York, 1979.
2. J. P. Aubin and I. Ekeland, "Applied Nonlinear Analysis," Wiley, New York, 1984.
3. H. Ben-El-Michaiekh and P. Deguire, Approachability and fixed point for non-convex set-valued maps, *J. Math. Anal. Appl.* **170** (1992), 477–500.
4. X. P. Ding and K. K. Tan, Generalized variational inequalities and generalized quasi-variational inequalities, *J. Math. Anal. Appl.* **148** (1990), 497–508.
5. K. Fan, A generalization of Tychonoff's fixed point theorem, *Math. Ann.* **142** (1961), 305–310.
6. K. Fan, Some properties of convex sets related to fixed point theorems, *Math. Ann.* **266** (1984), 519–537.
7. A. Granas and F. C. Liu, Coincidences for set-valued maps and minimax inequalities, *J. Math. Pures Appl.* **65** (1986), 119–148.
8. C. W. Ha, Minimax and fixed point theorems, *Math. Ann.* **248** (1980), 73–77.
9. B. Knaster, C. Kuratowski, and S. Mazurkiewicz, Ein Beweis des Fixpunktsatzes für n -dimensionale Simplexe, *Fund. Math.* **14** (1929), 132–137.
10. M. Lassonde, On the use of KKM multifunctions in fixed point theory and related topics, *J. Math. Anal. Appl.* **97** (1983), 151–201.
11. M. Lassonde, Fixed points for Kakutani factorizable multifunctions, *J. Math. Anal. Appl.* **152** (1990), 46–60.
12. S. Naoki, A further generalization of the Knaster-Kuratowski-Mazurkiewicz theorem, *Proc. Amer. Math. Soc.* **111** (1991), 187–195.
13. S. Park, Generalizations of Ky Fan's matching theorems and their applications, *J. Math. Anal. Appl.* **141** (1989), 164–176.
14. S. Park, Foundations of the KKM theory via coincidences of composites of upper semicontinuous maps, *J. Korean Math. Soc.* **31** (1994), 493–519.
15. S. Park, S. P. Singh and B. Watson, Some fixed point theorems for composites of acyclic maps, *Proc. Amer. Math. Soc.* **121** (1994), 1151–1158.
16. M. H. Shih and K. K. Tan, A geometric property of convex sets with applications to minimax type inequalities and fixed point theorems, *J. Austral. Math. Soc. Ser. A* **45** (1988), 169–183.
17. M. H. Shih and K. K. Tan, Generalized bi-quasi-variational inequalities, *J. Math. Anal. Appl.* **143** (1989), 66–85.
18. C. L. Yen, A minimax inequality and its applications to variational inequalities, *Pacific J. Math.* **97** (1981), 477–481.
19. E. Zeidler, "Nonlinear Functional Analysis and Its Applications," Vols. 1–5, Springer-Verlag, New York/Berlin, 1986–1991.